ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
10 CENTS PER COPY
SUBSCRIPTION PRICE, $4.00 PER YEAR

PURCHASER AGREES NOT TO RESELL OR DISTRIBUTE THIS
COPY FOR PROFIT.—PUB. RES. 57, APPROVED MAY 11, 1922

▽

JOURNAL OF AGRICULTURAL RESEARCH

VOL. XXV WASHINGTON, D. C., AUGUST 11, 1923 No. 6

CORRELATION OF FOLIAGE DEGENERATION DISEASES OF THE IRISH POTATO WITH VARIATIONS OF THE TUBER AND SPROUT [1]

By ALFRED H. GILBERT

Associate Pathologist, Vermont Agricultural Experiment Station

INTRODUCTION

It has been previously shown through the observations and experiments of a number of investigators that mosaic and leafroll of the Irish potato (*Solanum tuberosum*) are transmitted from season to season by means of the tubers. Among those who have contributed chiefly to this conclusion are, Orton (*9*),[2] Appel (*1*), Wortley (*15*), Stewart (*13*), Murphy (*8*), Melhus (*7*), Schultz et al. (*10*), and Schultz and Folsom (*12*). Since these diseases although carried in the tissues of the tubers do not cause any observable changes in their outward appearance, there would seem to be no means of distinguishing between healthy and diseased specimens when unsprouted tubers are held under favorable storage conditions. The observations and experiments reported in this paper were undertaken, therefore, with the purpose of discovering whether there were not some distinguishing characteristics associated either with the earlier or later stages of germination of tubers which might be utilized in determining the presence of mosaic or leafroll therein during the latter part of the storage period. It is evident that such correlations, if existent, would add to the knowledge of the diseases concerned by furnishing additional means of diagnosis and might also be a basis for the elimination of undesirable seed from stocks to be used for planting. The essential results of these investigations were reported in abstract at the Toronto meeting of the American Association for the Advancement of Science in December, 1921, and published in a subsequent number of Phytopathology (*6*).

Studies made at the Vermont Agricultural Experiment Station seem to indicate that certain definite correlations exist between both dormant and sprouted tubers and foliage condition in the plants grown from such tubers, and these studies were supplemented by observations both in the greenhouse and in the field as to variations in the seriousness of mosaic and leafroll symptoms in successive generations of plants and other related points.

In April, 1921, the noteworthy paper by Schultz and Folsom entitled "Leafroll, Net-Necrosis and Spindling-Sprout of the Irish Potato" (*12*) appeared, and though much wider in scope than the present paper and developed along different lines, yet certain of the most significant conclu-

[1] Accepted for publication May 2, 1923.

[2] Reference is made by number (italic) to "Literature cited," p. 265–266.

sions closely duplicate and corroborate those of the present investigation. This is notably true of the correlations shown between spindling-sprout and net-necrosis of the tuber and leafroll of the foliage. In addition to these correlations, the studies reported herein furnish evidence on other matters related to tuber germination and foliage symptoms—namely, (*a*) retardation in the sprouting of mosaic and leafroll tubers, (*b*) bud uniformity in mosaic and leafroll tubers, (*c*) the simultaneous occurrence of both mosaic and leafroll symptoms in the same plant, (*d*) variation in the progressive seriousness of mosaic infection from season to season, and (e) apical dominance in its relation to disease.

GERMINATION TESTS

During the winters of 1919–20 and 1920–21 many tubers of known history as to the condition of the plants from which they came were germinated. The germinations were made in the greenhouse under favorable conditions of moisture and temperature, the seed pieces being planted in soil in benches and pots or in sawdust and excelsior in wide shallow pots. In the course of the trials it was found that tubers from a crop planted in May and harvested in October did not germinate normally until the following January or February. In the length of the rest periods thus indicated the facts observed conform in general to the observations of Appleman (*2*), who found that the rest periods of Irish potatoes differed to some extent between varieties but were quite uniform within a certain variety. He found that while in general the rest period was terminated after about 90 days, it could be very much shortened by removing the skins from the tubers without injury to the buds. In the course of the investigations herein reported, earlier germination was secured by using Appleman's method, which, however, did not prove to be adapted to the needs of the present studies. The most satisfactory results were secured from germinations made toward the end of the storage season, during February, March, and April, while some of the most significant observations on spindling-sprout were made in the case of tubers sprouting in the storage cellar at the very end of the storage period. The cellar in which the tuber samples were kept during the winter was sufficiently moist so that the potatoes did not shrivel, and registered temperatures ranging from 35 to 45° F. throughout the winter months, rising with the higher temperatures of April and May. At planting time most of the tubers had developed sprouts of varying lengths.

RETARDATION IN SPROUTING

As a result of germinating many tubers both diseased and healthy in the greenhouse, evidence has accumulated that germination of the seed piece and growth of the young shoot are definitely and markedly retarded both in the case of serious mosaic and of serious leafroll infections. In instances of mild degrees of infection, however, little if any variation from the normal appearance in sprouting can be detected. In order to secure evidence on this point, tubers of the Green Mountain variety were used, some of them of known history as to the presence or absence of disease for one or more seasons, while others were derived from general stock. One half of each tuber was planted in a shallow pot on moist sawdust and covered with excelsior, and the other half was planted in soil in an 8-inch pot. All were given uniform treatment as to light, moisture, and temperature. The appearance and rate of

growth of the shoots from the diseased as compared with those from the healthy tubers were observably unlike. Slower growth occurred and a pinched, unhealthy and slightly abnormal appearance characterized the sprouts from tubers seriously affected with either mosaic or leafroll. Plate 1, A (two potted plants and sprouted seed pieces from the same corresponding tubers), illustrates this point. The plant at the left exhibits extreme mosaic symptoms and the sprouts of the seed piece distinct retardation in growth; the plant at the right, mild mosaic symptoms and a nearly normal sprout on the seed piece. Plate 1, B, shows another case of retarded sprouting, the potted plant being from the same tuber as the sprouted seed piece and exhibiting extreme mosaic symptoms. The plants shown in Plate 1, B, were from a parentage described as "healthy but adjacent to mosaic hills." Plate 1, A, shows an additional evidence of weakness and degeneration in the brown lesions on the several sprouts of the seed piece at the left. Schultz and Folsom (*11*, *p.* 321) have reported a "spotting and streaking of the leaves, petioles and stems" apparently somewhat similar to the foregoing, accompanying mosaic symptoms induced in potato plants by artificial inoculation.

RELATION OF MOSAIC AND LEAFROLL IN VERMONT

Schultz and Folsom (*12*, *p.* 75) note that in northeastern Maine the Irish Cobbler is the only variety which shows more than a trace of leafroll in commercial fields and that it "seldom shows more than a trace of mosaic, while the Green Mountain and Bliss Triumph varieties are seldom found with only a trace of mosaic."

In Vermont the facts with reference to these points are somewhat different. As in Maine, Irish Cobblers are more subject to leafroll than to mosaic, but this variety in Vermont frequently shows more than a trace of mosaic." Further, Green Mountains in Vermont are now quite as commonly, though in general not as seriously, affected with leafroll as with mosaic. The State certified seed-potato inspectors during the past two or three seasons have found fields with relatively high percentages of mosaic much more frequently than fields with high percentages of leafroll. It is also a common experience now in this State to find in many stocks of Green Mountains, and less frequently in Irish Cobblers, both mosaic and leafroll infection in the same field. The percentages of mosaic found are frequently reported as "only a trace." In these respects the conditions in Vermont with reference to mosaic and leafroll occurrence are appreciably different from those in Maine.

SIMULTANEOUS OCCURRENCE OF MOSAIC AND LEAFROLL IN THE SAME PLANTS

Stewart (*13*) in 1916, in recording his observations on "some degenerate strains of potatoes," reports that "certain hills were much affected with leafroll and showed also traces of mosaic." In studying the symptoms of mosaic and leafroll plants both in the field and in the greenhouse, it has frequently been observed by the writer that a plant may exhibit the characteristics of both diseases at the same time. Instances of this have been noted both in the early part of the season and later. When the symptoms appeared very early it was obvious that the double infection had been carried in the seed tuber, while later appearances might have been due to secondary infections of one or both diseases. In most cases where this double infection was observed, the unhealthy

condition of the plant was very marked. There was a spindling habit, foliage area was reduced, and in some instances brown necrotic regions had developed on leaves, petioles, and stems, and premature defoliation had occurred. Plate 1, C, shows a plant in which these symptoms were observed under greenhouse conditions.

In Table I it will be noted that several samples are recorded as exhibiting both mosaic and leafroll symptoms. In these cases the leafroll may perhaps be considered as the more significant since it is here associated with net-necrosis and spindling-sprout. The two infections combined seem to produce a condition of extreme weakness and low yield, more marked than would result, except occasionally, from either disease alone. The plants shown in Plate 6 also illustrate the double infection described. In these plants the symptoms of both foliage troubles were easily recognized. There is the mottling and wrinkling of leaflets, shortened petioles, and crowding of leaflets typical of mosaic, and, in addition, the rigid texture, the rolling of leaves, and the "staring" habit characteristic of leafroll.

Table I furnishes evidence on several points as follows: (*a*) The data tend to substantiate the statement made in another portion of this paper that spindling-sprout is correlated with leafroll but not with mosaic. It will be noted in this connection that mosaic symptoms are recorded as associated with leafroll in some of the plants from spindling-sprout tubers, but that no cases of leafroll are cited as occurring in the plants from normal sprouts. It is not to be assumed that symptoms of leafroll do not occur in plants from tubers with normal-appearing sprouts, but it is evident that there is a correlation between spindling-sprout and leafroll and, further, that this is independent of the mosaic symptoms. (*b*) Tubers with normal sprouts produced either mosaic or healthy plants and in general larger yields than tubers with spindling sprouts. The hill yields of the tuber units show a fairly uniform decrease, being largest in hills from bud-end seed pieces and smallest in hills from stem-end pieces. This was true not only of the leafroll and mosaic but of healthy tuber units.

TABLE I.—*Yields by tuber units from spindling-sprout and normally sprouted*

Record No. (1921).	Condition of sprouts.	Yields in grams of tuber unit hills—					Total.	Average per hill.	Condition of plant (1921).
		a.	b.	c.	d.	e.			
75 (16-1)	Spindling	184.1	116	67.5	93.7	45	506.3	101.2	Leafroll.
76 (16-2) [a]	do	224	182.8	194.5	124.5	256	982.4	196.4	Do.
77 (16-3)	do						231.7	77.2	Leafroll and mosaic.
82 (18-2)	do	111	67.3	39	40.5		258	64.5	Very weak leafroll and mosaic.
85 (19-2)	do	117	116				233	106.5	Leafroll and mosaic.
86 (19-3)	do	60	35	26.5	23.5		145	36.2	Leafroll.
87 [b]	do	87	42.5	31.4			159.9	53.3	Leafroll and mosaic.
89 [b]	do	139.9	79.4				219.3	109.6	Do.
78 (17-1)	Normal						659	164.7	Mosaic (medium).
79 (17-2)	do						760.8	253.6	Healthy.
80 (17-3)	do	573	421	362	175.7		1,471.7	367.9	Do.
81 (18-1)	do						958.7	191.7	Do.
83 (18-3)	do						232	46.4	Mosaic.
84 (19-1)	do	327	276.5	263	184.5		1,051	262.7	Do.
88	do	282	230	206	115		833.3	208.3	Healthy.
90	do	304.8	254				588.8	279.4	Mosaic.

[a] It is possible that an error in placing the seed pieces is responsible for the larger yield of hill *c* in this sample, since such a result is not in harmony with those from the other tuber units.

[b] Tubers showed net-necrosis when cut.

VARIATION IN THE PROGRESSIVE DEVELOPMENT OF MOSAIC SYMPTOMS

Schultz and Folsom (*11, p. 316*) recognize and describe five stages or degrees of severity of mosaic symptoms—"slight," "slight plus," "medium," "medium plus," and "bad." The writer, while agreeing fully as to the significance and appropriateness of these terms, has, however, used the slightly different terms, "mild," "medium," and "extreme," respectively, for the conditions described by Schultz and Folsom as "slight," "medium," and "bad." "Mild" mosaic in the present paper indicates faint mottling with approximately normal development of foliage; "medium" refers to a condition in which there is distinct mottling and some wrinkling and reduction in leaf area, and "extreme" is used in connection with the stage in which there is marked wrinkling and curling of leaves and a dwarfed or conspicuously spindling habit of the plant. The results of Schultz and Folsom (*11, p. 317*) indicate that mosaic in northern Maine "does not necessarily change much from year to year in any diseased stock after the first appearance of the effects of infection." The results of the present investigation in part seem to agree and in part to differ with the foregoing statement. That is to say, it has been found that in Vermont mild mosaic may in some cases persist as mild mosaic for several seasons in a certain stock without marked change, but that from one season to the next sudden changes also occur, as from mild to extreme, or even from a condition noted as "healthy but adjacent to mosaic hills" to a condition of extreme mosaic as described in the succeeding paragraphs.

As a result of observations on many plantings of tubers of known history both in the greenhouse and field, it was noted that much variation exists in the progressive development of mosaic infection from year to year. In certain cases a condition of mild mosaic persists in the progeny plants through one or more subsequent seasons with comparatively little change, while in other cases the increased seriousness of the disease as evidenced by the foliage symptoms is very marked between one season and the next. In the first instance tubers from plants exhibiting a mild degree of infection produced the next season plants showing little, if any, increase or change in the seriousness of the symptoms (Pl. 2, A 2; D), while in the latter case tubers from other plants with very similar external symptoms produced the next season plants showing, even in the earliest stages of development, extreme mosaic symptoms (Pl. 2, A 1; C).

In illustration of the first condition described is cited the history of some tubers of the Green Mountain variety secured in 1919 from L. P. Lloyd of Whitingham, Vt. The plants from which these tubers were taken were of large growth with leaves showing faint mottling but no wrinkling. There was apparently no reduction in area of leaf surface on account of the disease. Tubers from these hills planted in the experimental plots in 1920 gave plants showing only a faint mottling and no wrinkling or dwarfing throughout the season. Tubers from the 1920 plants in the field plots were in turn planted in the greenhouse in the winter of 1920–21, and the foliage produced showed faint mottling, slight wrinkling, and symptoms of leafroll. Since the plants in the field during the season of 1920 were adjacent to leafroll hills, there was sufficient opportunity for the acquisition of this infection. There is seen, then, in the foliage of three successive generations of this particular lot (III–18–2–1919) only a slight increase in the seriousness of mosaic symptoms,

and that in the third season. No record was obtainable of the disease status of this sample previous to 1919.

Referring now to the other condition mentioned previously—namely, that in which there seemed to occur a marked increase in seriousness of infection from one season to the next, the following cases are cited: (*a*) It was observed in plants just emerging from the soil in the greenhouse bench that certain ones exhibited marked wrinkling and mottling of leaves at this early stage (Pl. 2, B and C), while others, which later developed typical though mild mosaic symptoms, showed no mottling or wrinkling at the corresponding period of development (Pl. 2, D). The plants shown in Plate 2, C and D, are both from parentage indicated as mild mosaic but not from the same hill. (*b*) Again, a case of extreme mosaic is observed in a very young plant which is from a parentage indicated as "healthy but adjacent to mosaic hills" (Pl. 2, B).

In Plate 2 the potted plants A, 1 and A, 2 are grown from the same corresponding tubers as the plants shown in C and D. A, 1 and C from the same tuber developed early the extreme type of mosaic from mosaic mild ancestry, while A, 2 and D developed mild mosaic symptoms from an ancestry exhibiting the same type of infection.

The writer has at hand no definite evidence upon which to base an explanation of the differing types of mosaic infection just described. There may be some weight accorded to the suggestion that certain plants possess a factor of resistance to mosaic infection which is absent in other plants. It is also conceivable that the divergencies in the type and seriousness of symptoms observed may be due to differences in amount of infective material introduced by aphids or other agencies. Another and perhaps more plausible explanation is that in the plants exhibiting the mild and extreme symptoms, there may occur what are in effect distinct types or strains of mosaic possessing sufficient individuality to account for the persistence of either through successive generations of the host plant. The sudden development of the extreme type in such plants as are illustrated in Plate 2, B, and the continuation of the mild type in Plate 2, D, may be considered as furnishing some support for this argument. It is to be hoped that further investigations will throw light upon this point.

SPINDLING-SPROUT AND LEAFROLL

In 1916 Stewart (*13, p. 351*) in a discussion of the results of his "Observations on Some Degenerate Strains of Potatoes," summarizes his conclusions in part as follows: "Although spindling-sprout symptoms appeared occasionally in the tubers of plants affected with mosaic, leaf-roll, and curly-dwarf, they were too infrequent to warrant the conclusion that spindling-sprout is correlated with any of these three diseases." He further adds, "sprouting tests of tubers from leafroll, curly-dwarf, and mosaic plants show that spindling-sprout is not a symptom of these diseases." In general, the present investigation has shown that spindling-sprout, while it is not, as a rule, a symptom of curly-dwarf or mosaic, is a consistent symptom of leafroll and often also of net-necrosis. Stewart's procedure in noting the sprout condition of tubers from plants of certain observed foliage symptoms of the previous season, seems to the writer to be not so dependable with reference to the information sought as that which takes into account as well, and indeed, primarily, the foliage condition of the progeny plants. It must frequently occur

in plots or fields where both mosaic and leafroll are common that vines receive infection during the season which does not develop into external symptoms until the following season. For example, plants with apparently normal and healthy foliage throughout the season nevertheless produce tubers which in the following season develop plants with various stages of mosaic or leafroll. Evidently the sprouts of such tubers would have significance with reference to the foliage condition of the plants which they produced rather than of the plants from which they came. Further, in the case of net-necrosis of the tuber it has been shown by Schultz and Folsom (*12, p. 65–71*) that this condition suddenly appears and often as suddenly disappears so far as the symptoms are concerned, and that in its correlation with leafroll it depends upon "recency of infection." Consequently, with reference to this trouble also, the symptoms of the plants resulting from net-necrosis tubers are of more significance than those of the plants producing them.

Blodgett, Fernow, and Perry report in an abstract on "Testing Seed Potatoes for Mosaic and Leafroll" (5) that "tubers indexed as leafroll grew sprouts generally thinner than healthy potatoes but not always of the extreme spindling-sprout type," and state that by a preliminary sprouting method all but 3 per cent was removed from two lots containing 50 per cent of leafroll. These writers also secured partially successful results in climinating mosaic and leafroll from seed stock by growing one seed piece from each tuber as an early crop in the field previous to planting the main crop.

In the spring of 1921 about 60 hills were planted in the experimental plot with seed pieces from tubers which had sprouted in storage and whose sprouts showed a more or less spindling condition. Of these, 20 hills were of the Burbank variety and 40 hills were Green Mountain. The 20 hills of Burbank (No. 91, 1921) planted with seed pieces from tubers all of which had developed spindling sprouts showed 100 per cent characteristic leafroll. A control row of 26 hills (No. 92, 1921) of the same variety and lot but planted with tubers which had apparently normal sprouts, produced 92 per cent healthy hills and 8 per cent leafroll hills. Three of the leafroll hills from No. 91 (1921) and one healthy hill from No. 92, (1921) are shown in Plate 3.

The 40 hills of Green Mountains were planted in single tuber units with tubers selected from various samples of the preceding season, these tubers having developed in storage sprouts of more or less marked spindliness. Without exception, each hill of every tuber unit developed leafroll in varying degrees of seriousness. All the tuber units were planted in the same order, the seed pieces from the bud end or apex first and the others progressively in order to the stem end. With a large degree of uniformity the plants from the bud-end pieces were larger, more vigorous, and more productive, while a rather uniform decrease in size and productivity characterized the plants from the middle and stem-end pieces. Plate 4, A, B, C, and D, shows four of the five plants of tuber unit series 75 (16–1) illustrating characteristic leafroll symptoms and gradual decrease in size. The photographs of these hills were taken with the camera at the same distance in each case. All of the plants were rather small, the smallest being about 8 inches in height. In Plate 5, A, 75, may be seen the sprouted tuber from which these plants were grown. It will be observed that there is one rather vigorous sprout at the apex and that the remaining sprouts are more or less spindling, a condition very common in connection with this type of degeneracy. In

general, spindliness tends to begin in the sprouts at or near the stem end of the tuber, one or more sprouts at the apex often remaining normal in size and appearance except in the most extreme cases, such as is shown in Plate 5, D, 85, where all the sprouts have become spindling. That the varying degrees of spindliness of sprout are correlated with differences in vigor and productivity of the resulting plants is illustrated in Plate 4, A, B, C, D; Plate 6, A, B, C, D, E; and Tables I and II.

TABLE II.—*Yields in grams from average hills of Burbank plants which exhibited extreme and mild leafroll symptoms and normal foliage, respectively*

Record No. (1921).	Condition of sprout.	Yields from average hills—				Total.	Average per hill.	Condition of plants.
		1	2	3	4			
91a	Spindling	120.4	63.2	0	0	183.6	91.8	Extreme leafroll.
91b	...do...	329.9	195.9	0	0	525.8	262.9	Mild leafroll.
92	Normal	835.7	584	591	613	2,624.5	656.1	Healthy.

In Table II, No. 91a and 91b included tubers selected for very much marked spindliness of sprout. As indicated previously, all the plants from these tubers developed characteristic leafroll. The leafroll symptoms, however, varied in seriousness and the plants showing extreme symptoms, being much dwarfed, were indicated as 91a, while those of larger growth but still with well-marked leafroll characters, were designated as 91b. Average hills from each of these groups were selected and the yields weighed, with the results given in the table. For comparison, four average healthy hills from lot 92 (1921) were selected and the yields weighed. The results correspond closely to expectation as governed by the condition of the foliage. Wide variation in yields between the two types of leafroll appears, and a difference nearly as great between mild leafroll and healthy hills.

CORRELATION OF NET-NECROSIS WITH SPINDLING-SPROUT AND LEAFROLL

In cutting the tubers of lot 91 (1921), referred to in preceding paragraphs, it was noted that every tuber showed typical net-necrosis of the sort illustrated in Plate 6, A, affecting a portion of its tissues. These tubers had been selected for investigation because of the extreme spindliness of the sprouts which had developed in the storage cellar during early spring. In the experimental plot the plants from these tubers developed 100 per cent leafroll. Many of the hills were very weak and some seed pieces produced no plants at all. In lot 92 (1921) of the same variety selected for normal sprouts no net-necrosis was found, and of this lot, as noted previously, 92 per cent of the plants were healthy and 8 per cent showed leafroll. A tuber unit series from a net-necrosis tuber is illustrated in Plate 6. Three of the four seed pieces of this tuber showed necrosis of the cortical region and from near the edges of the necrotic tissue in two of the pieces sprouts which were thin and spindling had developed. The seed piece from the apex of the tuber developed two sprouts apparently normal, but the plant from this piece, as well as those from the pieces showing necrosis, developed both leafroll and mosaic symptoms. The occurrence of the mosaic symptoms in this instance, as in some others observed, is be-

lieved to be incidental so far as its connection with net-necrosis is concerned. A study of Table I will reveal the fact that leafroll resulted in every case of spindling-sprout, either alone or associated with mosaic, while there was no case of spindling-sprout observed when mosaic alone appeared in the resulting plants. On the other hand, numerous cases of mosaic of both the mild and extreme types were recorded in plants from tubers with normal sprouts. Similar conclusions are reached with reference to net-necrosis—that is, that while both mosaic and leafroll symptoms have been noted in plants from net-necrosis tubers, in no cases have such tubers resulted in plants exhibiting mosaic symptoms only. Net-necrosis, therefore, of the type discussed and illustrated in the present paper, is definitely and closely correlated with spindling-sprout of the tuber and with leafroll of the plant.

TABLE III.—*Showing correlation of spindling-sprout and net-necrosis of the tuber with leafroll of the foliage*

Lot No.	Number of tubers.	Condition of sprouts.	Net-necrosis.	Number of hills.	Condition of foliage.
..	1	Spindling....	Present.....	4	Leafroll and mosaic.
..	1	do.....	do.....	3	Do.
..	8	do.....	do.....	20	Leafroll.
..	1	do.....	do.....	4	Do.
..	3	do.....	do.....	13	Do.
..	2	do.....	do.....	4	Do.

APICAL DOMINANCE

Appleman in an article entitled "Some Factors Influencing the Vitality of Seed Potatoes" (*4*) in Maryland Farmer, volume 5, No. 16, 1921, states that "good normal tubers will as a rule sprout first from the seed or apical end," and also that "lack of apical dominance is a sign of low vitality in the tuber or in other words of its inability to produce vigorous plants." In another publication (*3, p. 80*) this author says: "The greater vigor of the terminal shoot may be due to greater inherent strength of the terminal bud or it may be merely apparent because the terminal shoot inhibits the growth of the lateral ones. The latter view suits best the requirements regarding the growth of buds on the potato tuber." He further states that while, under uniform conditions, terminal buds ordinarily grow out first, still if the basal eyes are more favorably situated with respect to the action of external agents the suppression of their growth by terminal buds may be entirely overcome. These facts are understood to be true only of the first crop of sprouts from a tuber.

In connection with the studies reported in the present paper many observations have been made as to the manner of sprouting of diseased and healthy tubers ,some of which have a bearing upon the matter of apical dominance. Tubers with normal sprouts from the apex and with the other eyes dormant were selected for planting and also those with normal sprouts from various eyes scattered over the tubers. Table IV gives the results of the observations upon the plants grown from these tubers.

TABLE IV.—*Showing relation between distribution of sprouts and foliage condition of progeny plants*

Lot No.	Number of tubers.	Distribution of sprouts.	Number of plants.	Condition of plants.
61	1	Clustered at apex	4	Medium mosaic.
62	1	do	4	Healthy.
63	1	do	2	Do.
65	1	do	4	Extreme mosaic.
66	1	do	3	Do.
67	1	do	4	Do.
78	1	do	4	Do.
79	1	do	3	Healthy.
80	1	do	4	Do.
57	1	Scattered	6	Medium mosaic.
58	1	do	5	Mild mosaic.
59	1	do	3	Medium mosaic.
60	1	do	3	Extreme mosaic.
64	1	do	5	Mild mosaic.
68	1	do	6	Medium mosaic.
69	1	do	5	Healthy.
70	1	do	4	Mild mosaic.
81	1	do	8	Healthy.
83	1	do	5	Mild mosaic.
84	1	do	4	Healthy.
88	1	do	4	Medium mosaic.
90	1	do	2	Extreme mosaic.

The results of these limited observations do not show either apical dominance or the lack of it in the tubers to be consistently correlated with any condition as to health or disease in the resulting plants. Tubers with sprouts clustered at the apex produced both healthy and mosaic plants and the same results were observed in the case of tubers with sprouts not confined to the bud end or apex. In Plate 5, B, 78, 79, 80 are shown tubers with normal sprouts clustered at the apex, and in C, 81, 83, and D, 84, of the same plate are shown tubers without this apical dominance. By referring to Table I it will be seen that of the tubers showing apical dominance, No. 78 produced mosaic plants, while No. 79 and 80 gave healthy plants, also, that of the tubers not exhibiting this dominance, No. 81 gave healthy plants while No. 83 and 84 gave mosaic plants.

In agreement with the statement of Appleman referred to above, the present observations indicate that apical dominance may be suppressed when basal eyes are situated more favorably for germination than terminal ones, so as an index of vigor in tubers this characteristic would be subject to limitations. Moreover, as an index of vigor it is subject to the further limitation that tubers with distinct apical dominance may develop mosaic and even mosaic of the extreme type, as in tubers nos. 66, 67, and 78 in Table IV.

Referring again to the data in Table IV, the records show that in the germination of tubers 61, 62, and 63 the plants from bud-end seed pieces started more promptly and grew better than the plants from the stem end of the tubers. The records further show in a limited number of cases examined that, regardless of disease condition, when there is marked apical dominance of sprouts, the plants from the apex grow more vigorously and those from the stem end less so, while in cases of

tubers with scattered sprouts, all apparently of equal vigor, the plants from the several seed pieces are approximately uniform. It seems to be true that a degree of apical dominance is shown in the material studied, in the large size and yields of hills from apical buds. The fact that should be emphasized, however, is that this dominance is not an index of freedom either from mosaic or leafroll.

The writer's conclusion with reference to the bearing of apical dominance upon the present problem is that the appearance of the sprouts as to normal thickness, branching, etc., is a better index of the healthy or diseased condition of the tuber and of the resulting plants than the distribution of the sprouts on the tuber; and further, that the data so far collected fails to support any consistent correlation between the distribution of sprouts on the tubers in storage and mosaic of the resulting plants.

SUMMARY

(1) In instances both of serious mosaic and serious leafroll infection, germination of the seed pieces and growth of the shoots are markedly retarded.

(2) (*a*) In the strains of Green Mountain potatoes under observation in these investigations the same plants frequently exhibited both mosaic and leafroll infection.

(*b*) All the data secured support the thesis that there is uniformity as to condition of health or disease in all the plants from a single tuber and also in all the tubers from a single hill.

(3) Marked variation in the progressive development of mosaic infection with the suggestion of differing strains of mosaic virus were indicated in the germination of tubers of known ancestry and in the observation of several successive generations of plants from known sources.

(4) (*a*) Spindling-sprout of the tubers is shown to be, in the varieties of potatoes studied, a consistent symptom of leafroll, but not of mosaic.

(*b*) Net-necrosis, of the phloem-necrosis type, is correlated in the tubers with spindliness of sprout and seems to be a consistent symptom of leafroll. The necrosis symptoms are not persistent in the progeny tubers.

(5) Yields of plants from spindling-sprout tubers were much reduced in comparison with those of plants from tubers with normal sprouts, and hills exhibiting extreme leafroll yielded far less than those with mild leafroll symptoms.

(6) Apical dominance in sprouting which has been held to be an indication of vigor in tubers was found to be associated both with the development of mosaic in the foliage and with a healthy condition of plants, and conversely, the lack of apical dominance was associated with the production of healthy as well as mosaic hills.

LITERATURE CITED

APPEL, Otto.

1915. LEAF ROLL DISEASES OF THE POTATO. *In* Phytopathology, v. 5, p. 139–148.

APPLEMAN, Charles O.

1914. BIOCHEMICAL AND PHYSIOLOGICAL STUDY OF THE REST PERIOD IN THE TUBERS OF SOLANUM TUBEROSUM. Md. Agr. Exp. Sta. Bul. 183, p. 181–226, 17 fig. Literature cited, p. 225–226.

1918. PHYSIOLOGICAL BASIS FOR THE PREPARATION OF POTATOES FOR SEED. Md. Agr. Exp. Sta. Bul. 212, p. 79–102, 11 fig.

(4) APPLEMAN, CHARLES O.
1921. SOME FACTORS INFLUENCING THE VITALITY OF SEED POTATOES. *In* Maryland Farmer, v. 5, no. 16, p. 5.

(5) BLODGETT, F. M., FERNOW, Karl, and PERRY, F. R.
1922. TESTING SEED POTATOES FOR MOSAIC AND LEAF ROLL. II. (Abstract.) *In* Phytopathology, v. 12, p. 40.

(6) GILBERT, A. H.
1922. THE CORRELATION OF FOLIAGE-DEGENERATION DISEASES OF THE IRISH POTATO WITH VARIATIONS OF THE TUBER AND SPROUT. (Abstract.) *In* Phytopathology, v. 12, p. 40.

(7) MELHUS, I. E.
1917. NOTES ON MOSAIC SYMPTOMS OF IRISH POTATOES. (Abstract.) *In* Phytopathology, v. 7, p. 71.

(8) MURPHY, Paul A.
1917. THE MOSAIC DISEASE OF POTATOES. *In* Agr. Gazette Canada, v. 4, p. 345–349, illus.

(9) ORTON, W. A.
1914. POTATO WILT, LEAF-ROLL, AND RELATED DISEASES. U. S. Dept. Agr. Bul. 64, 48 p., 16 pl. Bibliography, p. 44–48.

(10) SCHULTZ, E. S., and others.
1919. INVESTIGATIONS ON THE MOSAIC DISEASE OF THE IRISH POTATO. *In* Jour. Agr. Research, v. 17, p. 247–274, pl. A–B (col.), 25–30. Literature cited, p. 272–273.

(11) ——— and FOLSOM, Donald.
1920. TRANSMISSION OF THE MOSAIC DISEASE OF IRISH POTATOES. *In* Jour. Agr. Research, v. 19, p. 315–338, pl. 49–56.

(12) ——— ——— ———
1921. LEAFROLL, NET-NECROSIS, AND SPINDLING-SPROUT OF THE IRISH POTATO. *In* Jour. Agr. Research, v. 21, p. 47–80, 12 pl. Literature cited, p. 78–80.

(13) STEWART, F. C.
1916. OBSERVATIONS ON SOME DEGENERATE STRAINS OF POTATOES. N. Y. (State) Agr. Exp. Sta. Bul. 422, p. 319–357, 12 pl.

(14) ——— and SIRRINE, F. A.
1915. THE SPINDLING-SPROUT DISEASE OF POTATOES. N. Y. (State) Agr. Exp. Sta. Bul. 399, p. 133–143, 3 pl.

(15) WORTLEY, E. J.
1915. THE TRANSMISSION OF POTATO MOSAIC THROUGH THE TUBER. *In* Science, n. s., v. 42, p. 460–461.

(16) ———
1918. POTATO LEAFROLL: ITS DIAGNOSIS AND CAUSE. *In* Phytopathology v. 8, p. 507–529, 16 fig. Literature cited, p. 529.

PLATE 1

A.—Two potted plants of the Green Mountain variety and sprouted seed pieces from the same corresponding tubers. On the left the sprouts of the seed piece are retarded in growth and show several brownish lesions, and the potted plant exhibits symptoms of extreme mosaic; on the right the seed piece has a sprout approximately normal, and the plant exhibits mild mosaic.

B.—A potted plant with symptoms of extreme mosaic and sprouted seed piece from the same tuber. The sprout exhibits retarded growth though not as marked as in A. The seed tuber was from a parentage described as "healthy but adjacent to mosaic plants."

C. A potted plant of the Green Mountain variety exhibiting symptoms of both leafroll and mosaic. Irregular brownish areas appeared in the leaves and on the stems, and premature defoliation occurred.


A


B
C



A
1
2
B
C
D

PLATE 2

A.—Plants of the Green Mountain variety exhibiting two types of mosaic symptoms, grown in greenhouse. 1, showing marked spindling habit and reduced leaf surface accompanied by wrinkling and mottling, the extreme type. 2, foliage nearly normal as to development but with the characteristic mottling, the mild type. Both plants are from mild mosaic parentage but from different hills.

B. Young Green Mountain plant showing early the symptoms of extreme mosaic, grown in greenhouse bench (1920–21) from parentage designated as "healthy but adjacent to mosaic hills," in the experimental field in 1920.

C. Young Green Mountain plant of the extreme type, developing the symptoms very early in the greenhouse bench (1920–21), from parentage designated as mild mosaic in the experimental field in 1920. From the same tuber as A, 1.

D.—Young plant apparently healthy at first but later developing mild mosaic symptoms. From mild mosaic parentage and from the same seed tuber as A, 2.


PLATE 3

A.—Healthy hill of Burbank variety (lot No. 92) grown from tuber with normal sprouts.

B, C, D.—Leafroll hills of the same variety as A but grown from tubers with spindling sprouts and net necrosis (lot No. 91). All plants of this lot developed characteristic leafroll.

Potato Foliage Degeneration Diseases

PLATE 3



Agricultural Research

Washington, D. C.



A
B
C
D

PLATE 4

Four hills from tuber No. 75 (16–1) planted in the order A, B, C, D, with seed pieces taken as follows:

A.—From the bud end.

B, C.—From the middle.

D.—From the stem end of the tuber.

All hills exhibit characteristic leafroll. The seed tuber developed spindling sprouts from several eyes. See plate 5, A.

54484—23——2

PLATE 5


Green Mountain tubers which sprouted in the storage cellar in the spring.

A.—Tubers 75, 76, and 77 illustrate a condition, frequently observed, in which the is one apparently normal sprout at the bud end, the remaining sprouts being more less spindling. All hills from these tubers showed leafroll or combined leafroll mosaic symptoms.

B.—Tubers 78, 79, and 80 with normal sprouts clustered at the bud ends. Tuber produced mosaic plants while 79 and 80 gave plants entirely healthy throughout season.

C.—Tuber 81 produced healthy plants; 82, very weak plants showing both leaf and mosaic (note the spindling sprouts toward the stem end); and 83, mosaic plan

D.—Tuber 84 with normal sprouts produced medium mosaic hills, while tubers and 86, with spindling sprouts, produced very weak plants showing leafroll sympto


A
77
76
75
B
80
78
79
C
82
83
81
D
86
85
84



A
2
3
4
B
C
D
E

PLATE 6

A. Seed pieces of tuber No. 87 exhibiting spindliness of some of the sprouts and necrosis of the tuber.

B, C, D, and E.—Plants grown from seed pieces 1, 2, 3, and 4, respectively. Both leaf roll and mosaic symptoms developed in these plants. A gradual decrease in size and vigor is noted in the plants from the middle and stem-end pieces as compared with that from the bud end. The bud-end piece showed neither necrosis nor spindliness of sprout, but produced, like the others, a diseased plant.

COMPARATIVE STUDY OF PHYTOPHTHORA FABERI ON COCONUT AND CACAO IN THE PHILIPPINE ISLANDS.[1]

OTTO AUGUST REINKING[2]

Formerly Collaborator, Office of Crop Physiology and Breeding Investigations, Bureau of Plant Industry, United States Department of Agriculture; Plant Pathologist Philippine Agricultural Experiment Station; and Professor of Plant Pathology, College of Agriculture, University of the Philippines

INTRODUCTION

Each year in the Philippine Islands, severe losses result from coconut budrot and from blackrot and canker of cacao. In 1919, the writer (7)[3] produced evidence indicating that the organisms causing both diseases were identical. These studies have been continued, and detailed cross inoculation experiments, comparisions of parallel cultures, and more complete physiological and morphological studies have been made.

Coconut budrot is a serious disease of the coconut (*Cocos nucifera*), affecting the single terminal bud of the tree. Throughout the world various organisms, including bacteria and fungi, have been considered responsible for the disease. In the Philippine Islands there are apparently two types of budrot—the really infectious type caused by a Phytophthora, and a secondary type following some injury such as beetle injury and caused primarily by the invasion of bacteria in the weakened tissue. The earliest symptoms of the infectious type are usually the production of rows of dark-brown spots on the newly unfolded leaves. Later developed leaves then show a severe leaf blighting. Finally, the central group of folded leaves dies and turns brown, indicating the death of the growing point. The bud may not be killed until after 12 months from the time of the first leaf spotting. The usual period, however, is from 2 to 3 months. The fringe of older, healthy, green leaves remains on the tree until it dies a natural death. Infected trees are stunted (Pl. 4, B) and the nuts do not develop. Internal symptoms are characterized by a browning of the folded leaves above the growing point. A white mycelial felt may or may not be present on these young leaves. From the growing point, downward and to the sides, the organism advances for a short distance into the woody parts, where the limits of its advance are marked by a dark red to brownish line (Pl. 1, A). In advanced cases, after the entrance of bacteria, the soft region about the growing point is changed to a semiliquid, ill-smelling mass.

The blackrot of pods and the canker of cacao (*Theobroma cacao*) is widely distributed throughout the world in the cacao growing regions. In the Philippines, it has been reported as causing severe losses (6, *p.* 192–196). The fruit is attacked at any stage of its development. Usually the greatest damage is done to the young fruits. At first a minute black spot is developed on the infected part. This spot gradually

[1] Accepted for publication May 2, 1923. Presented to the graduate school of the University of Wisconsin in partial fulfillment of the requirements for the degree of doctor of philosophy.

[2] Plant pathologist of the United Fruit Co. in Honduras and Guatemala.

[3] Reference is made by number (italic) to "Literature cited," p. 284.

Journal of Agricultural Research, Washington, D. C.

Vol. XXV, No. 6
Aug. 11, 1923.
Key No. G-324

enlarges until the pod becomes blackened. In excessive damp weather a mass of mycelium with conidia may be produced on the surface of diseased fruit (Pl. 2, B). The mycelium invades the rind, passing into the seed. Diseased pods may fall or remain on the tree, drying up and becoming mummified. Flowers and stems may be attacked and killed by the fungus. The canker is found on young twigs, older branches, and the trunks of trees. Diseased twigs are characterized by browning of the leaves, dying of the tips, and shrivelling of the diseased wood. The first evidence of a canker is the appearance of a darker than normal color on the bark. The infected area may then turn brownish. A shrinking of the diseased area and a definite line of demarcation is usually produced, showing the limits of the infection. In advanced stages the bark may exude an amber fluid and later may crack and scale. The infection has been observed to spread from diseased pods into the branches or trunk. A true cankered condition is not always produced. Internal symptoms are characterized by a browning and blackening of the infected bark and wood (Pl. 2, A).

The fungus from coconut, used for the comparative study, was isolated from an infected coconut tree on March 11, 1919. The diseased tree was located in a plantation near Lilio, Laguna Province, P. I., a region in which sporadic epidemics of the disease have been recorded since 1905.

The fungus from cacao was isolated from a diseased cacao pod collected in the plantings near Los Banos, Laguna Province, P. I., during 1917. The disease is present there in a severe form throughout the year, causing most damage during the rainy season.

COMPARATIVE STUDY OF THE FUNGUS ON BOTH HOSTS

FIELD CHARACTERISTICS

On coconuts the single terminal bud of the tree is affected. A white mycelial felt may or may not be present on the young, folded, diseased leaves of the bud. Microscopic examination of the infected parts of the leaves, growing point, and surrounding tissue shows the presence of a granular, nonseptate mycelium in the intercellular spaces (Pl. 8, A). Fingerlike haustoria are produced in abundance, penetrating the host cells (Pl. 8, B). Chlamydospores, in limited numbers, have been observed in the infected buds some 60 cm. above the growing point. The writer has never observed the lateral penetration into the tender bud through the leaf bases. The penetration in all cases observed appeared to be a vertical, downward one, along the central leaf sheaths into the bud.

Diseased cacao pods may be covered with a mass of mycelium with conidia, but usually there is no macroscopic evidence of mycelium (Pl. 2, B). Conidia are produced in abundance on the surface of the fruit. Microscopic examination of the interior of diseased pods shows the presence of a granular, nonseptate, intercellular mycelium and usually a mass of chlamydospores (Pl. 8, C). In cankered areas the mycelium develops primarily internally in the diseased tissue. Rarely, as under excessive damp conditions, the hyphae are produced on the surface.

INFECTION EXPERIMENTS

The evidence given before by the writer (7) is sufficient to prove that the organism from coconut budrot can produce the disease in coconut seedlings and older trees through injuries. Further infection experi-

ments have been made with both the stab method of inoculation and inoculation of uninjured trees. In practically every instance disease was produced. Out of 17 seedlings inoculated by the stab method, 16 were severely infected (Pl. 5). Of 7 trees inoculated by merely pouring a zoospore suspension of the fungus into the bud, 6 contracted the disease and were killed. All controls remained healthy, and reisolations of the fungus were readily made from the infected trees.

One special case of inoculation of a healthy, vigorous tree under control conditions in the field bears further mention. The three trees used for this test were situated on the Faculty Hill of the College of Agriculture at Los Banos, Laguna Province, P. I. The trees, located near the writer's home, were far removed from other coconuts and were extremely healthy, being free from serious fungus attacks and insect infestation. They grew on a hillside where there was a good air movement and, consequently, a location not peculiarly adapted for the best development of the disease. The trees were approximately 3 years old at the time of inoculation. Cultures 1 month and 11 days old growing on corn meal were used for the inoculations. On November 20, 1919, rain water was poured into one culture flask and then as soon as zoospores were produced, the zoospore suspension was poured between the young folded leaves of the bud. Another tree was similarly inoculated, except that distilled water was used in place of the rain water. For a control the third tree was used. An equal amount of rain water free from the organism was poured into the bud. On December 19, 1919, the tree inoculated with the rain water suspension showed evidences of disease. The newly unfolded leaves had a series of brown spots on the leaflets forming a concentric ring across the entire leaf. The bud from all outward appearances was entirely healthy. This spotting continued to be produced on the newly developed leaves, becoming gradually more pronounced. On April 16, 1920, the spotting was severe. August 6, 1920, the bud apparently was still healthy, but the newly unfolded leaves were badly spotted. A severe leaf blight was observed on October 19, 1920, and the tree showed evidence of being stunted. The bud appeared to be somewhat weakened. On January 8, 1921, the bud was almost dead, being browned, and on January 25, 1921, it was dead. The other two trees were healthy and showed a marked increase in growth over the diseased tree (Pl. 4, B). A typical case of budrot was produced in the infected tree (Pl. 3, A). Under field conditions in localities where epidemics of the disease are severe, the trees succumb faster, usually within two or three months from the time of infection. The inoculated tree was finally cut down and a longitudinal section through the bud and growing point showed the disease to be typical of the naturally infected cases of budrot (Pl. 4, A). The control, as well as the tree inoculated with a zoospore suspension in distilled water, remained healthy.

Besides the coconut palm (*Cocos nucifera*), other members of the Palmae were also successfully inoculated with the organism, and disease was produced in a severe or medium form. The stab method of inoculation was used. A stab with a sterilized scalpel was made through or near the growing point and then the fungus was inserted. In each case the fungus was recovered from the diseased trees. The controls all remained healthy.

The following seedlings were killed by the fungus attack: *Archontophœnix alixandraeae, Dypsis madagascariensis, Livistona rotundifolia, Normanbya merrillie. Pinanga insignis.*

Livistona rotundifolia appeared to be the most susceptible.

A medium to severe infection was produced in the seedlings *Phoenix dactylifera* and *Ptychosperma mearthurii*.

Evidence previously given (7) has also proved conclusively that the Phytophthora isolated from cacao will produce a typical case of coconut budrot in injured and uninjured trees (Pl. 3, B). The fungus isolated from cacao and used for the comparative tests was also highly parasitic on the pods and branches of cacao (Pl. 2).

In order to determine whether the two fungus strains would both, in a similar manner, attack a variety of hosts, various inoculation experiments were conducted. It is realized that these inoculation tests can not alone be used to distinguish between various species of Phytophthora, but taken into consideration with the physiological and morphological characters, it is an added step toward proving the similarity or dissimilarity of two of the strains. In every instance in the inoculation experiments, reisolations of the fungus from the diseased plants were successfully made. All controls remained healthy. The results of the various cross inoculations in which cases both strains of Phytophthora, one from coconut budrot and the other from blackrot of cacao, were used, are summarized as follows:

Seedlings severely infected by both strains—
Annona muricata (guanabanos, soursop).
Cocos nucifera (coconut).
Hevea brasiliensis (Hevea or Para rubber).
Theobroma cacao (cacao).

Seedlings slightly infected by both strains—
Annona glabra (annonas).
Mangifera indica (mango).
Sandoricum Koetjape (santol).

Fruitrot produced by both strains—
Carica papaya (papaya).
Lycopersicum esculentum (tomato).
Malus malus (apple).
Theobroma cacao (cacao).

Tuber-rot produced by both strains—
Solanum tuberosum (Irish potato).

Vegetable blight produced by both strains—
Pisum sativum (pea).

The coconut and cacao seedlings were readily killed by inoculation with either strain of the fungus. A severe blight of the Hevea rubber seedlings was produced. Under certain conditions a stemrot of 6 cm. developed in three days. On the soursop a rot 5 to 6 cm. in extent was formed by the attack of both strains.

The rotting of papaya fruit was brought about in the same manner by both. After the first day, the fungus mycelium was well developed on the surface and in the interior of the fruit. In the early stages a white, cottonlike growth was formed (Pl. 1, B). The diseased part was slightly discolored, being a darker yellow, especially marked at the boundary of the circular advance of the fungus. Microscopic examination showed the presence of the mycelium within the diseased tissue and the production of numerous spores on the surface. A softrot was finally formed.

Ripe and green tomato fruits were also rotted by both strains of the fungus. The appearance and rapidity of the rot was identical for both

strains. The green fruits were not so rapidly attacked. In two days there was only a slight rot produced on the green fruits, but a severe rot on the ripe ones. Later a dense growth of mycelium developed on the surface, appearing first on the ripe fruits. At the end of nine days, all fruits, both green and ripe, were severely rotted and covered with a mass of white mycelium which arose through the epidermis from within the fruit (Pl. 6, A and B). The green fruits were browned and darkened within, while the ripe fruits were only softened. Microscopic examination showed the presence of an intercellular mycelium throughout the entire fruits. A pure culture of the fungus was present on the surface with many chlamydospores and conidia.

Ben Davis apples were easily attacked and rotted by both strains of the fungus. Within five days after the inoculation through the epidermis, a rot spreading 4 cm. beneath the skin and extending to the core was produced (Pl. 6, D and E). The rotted area was brown and rather mealy. Microscopic examination showed an abundance of characteristic intercellular mycelium throughout the rotted area.

Both strains of the fungus infected cacao pods, developing the characteristic blackrot. The strain isolated from cacao appeared to be more virulent.

The rot of potatoes was produced rather slowly by both strains. Irish Cobbler and Rural New Yorker potatoes were used for the inoculations. After 6 days a slight rot developed and within 30 days the rot extended to the center of the tubers. The infected potatoes, in certain cases, showed the symptoms as recorded by Pethybridge (5) on the rotting of potato tubers by a new species of Phytophthora. No definite line of demarcation between the healthy and diseased portions was noted immediately after cutting the tubers. A blackish line at the extremity of the infection and following around with the vascular system just beneath the surface then developed. Soon after cutting the tubers, the affected portion appeared somewhat watery, and in 15 to 30 minutes a distinct pinkish to red coloration was formed in the rotted region. Later, on longer exposure to the air, the invaded parts turned a purplish black. Not in all cases of infection did the pinkish or reddish coloration develop. Microscopic examination of the infected areas showed the presence of an intercellular, nonseptate, granular mycelium.

Garden bush pea plants were readily attacked and severely blighted by both strains of the fungus. In six days after inoculation a severe blight was produced. A whitish mass of mycelium formed over the blighted parts. The mycelium was present in the intercellular spaces of the affected tissue.

The consistent similarity in the attack of various hosts by both strains of the fungus shows that in this respect there is a constant likeness. In no case, except probably with the cacao fruit, was there any difference noted in the virulence of the two. Both are omnivorous, capable of attacking a large number of different hosts.

PHYSIOLOGICAL CHARACTERISTICS

GROWTH ON VARIOUS MEDIA

The two strains grew well on a variety of media, the character of the growth being similar for each. Since the two grew alike in all particulars, a separate description of each will not be given. The following discussion applies equally well to either. The descriptions were made

from cultures growing in the ordinary laboratory light under tropical room temperatures.

CORN MEAL.—At first the growth on corn meal was very scant, being hardly recognized. The mycelium spread slowly and thinly in a circular area over the medium. Few, short, aerial hyphae were produced. In two to four days the growth was sparse and granular, due to the production of spores. As the culture grew older, the hyphae became more in evidence and a distinct granular appearance was formed. Finally a slight cottony, much granular growth developed over the surface of the corn meal. An abundance of conidia and fewer chlamydospores were produced.

LIMA BEAN AGAR (100 gm. lima beans, 20 gm. agar, 1,000 cc. water).—Growth started with few aerial hyphae that gradually spread over the surface of the agar slant. In three days the mycelium had nearly spread over the surface. It was then characterized by being aerial, white, cottony, and granular. The granular appearance, due to spore production, was more in evidence on the walls of the glass tube at the edges of the growth. As the culture grew older, there was a gradual thickening of the aerial, mycelium mass.

OAT MEAL AGAR (100 gm. Quaker Oats, 17 gm. agar, 1,000 cc. water).—At first a few aerial hyphae developed which then gradually spread over the surface of the slant. After four days the mycelium, although still not very thick, had spread over the whole surface of the agar. Many granular bodies developed at this stage. The growth was exceedingly rapid and soon the mycelium formed a thick mass evenly spread over the agar slant, being slightly thicker at the base. As the culture grew older the hyphal mass gradually became more and more dense. Chlamydospores were produced in abundance and conidia were not so much in evidence.

POTATO DEXTROSE AGAR (100 gm. potatoes, 20 gm. dextrose, 15 gm. agar, 1,000 cc. water).—The first growth was characterized by being flocculent and submerged in the medium. On the second day, aerial hyphae appeared along with an increase of the submerged, flocculent growth. The aerial hyphae gradually became more dense and covered the entire slant. After five days the vegetation was dense, white, cottony, and usually thicker on the lower part of the slant. Minute granules formed by the spores were soon in evidence. Chlamydospores appeared to be produced first and in more abundance than the conidia.

RELATION TO LIGHT

In all of the cultures after exposure to the ordinary laboratory light for four days spores were produced copiously. If the cultures were kept in a dark chamber or incubator, free from light, only a few spores were formed. Sporulation was therefore hindered to a certain extent by darkness. There were apparently more chlamydospores than conidia produced in the dark. The growth in the light was more granular than that in the dark.

RELATION TO TEMPERATURE

Both strains of the fungus behaved the same in their relation to temperature. Altmann controlled temperature incubators ranging from about 4° to 32° were used for these determinations. A parallel series on potato dextrose agar plates was placed at the various temperatures and the difference in vegetative growth was determined by measure-

ments of the diameter of the mycelial mass in each case. Very slight growth for both strains was noted at 12°. The vegetation gradually increased in extent up to between 27° and 30°. From 30° and above there was a gradual decrease. The optimum growth occurred between 27° and 30°, being nearer 27°. Growth still was very good at a temperature of 32°. Below 12° no growth was noted. Neither fungus strain, however, was killed when held at an average temperature of 11.5° for 16 days, for growth took place after removal from the incubator. When kept at an average temperature of 7.5°, the minimum being about 6° for the same period, they were killed.

In all cases of growth the hyphae grew into the agar. At the lower temperatures the mycelium formed a dense, flat, surface mat with few aerial hyphae. At temperatures of 22°, and above, a tufted, serial mass of hyphae was produced, radiating out from the center. No spores were formed at the lower temperatures, between 12° and 20°. Few conidia and chlamydospores developed at the higher temperatures on the plates kept in the dark. In cases where spores formed in the dark, the chlamydospores appeared to predominate. There was a marked contrast between the growth on plates of both strains of the fungus kept in the light at a room temperature of about 22° and that on plates kept in the dark in the incubator at the same temperature. In both cases the vegetative development was approximately the same in extent. In the dark, however, the granular development of aerial hyphae was not produced and few conidia and chlamydospores formed. The growth on plates in the light was tufted, aerial, and granular, with an abundance of both chlamydospores and conidia. The development of the coconut strain is clearly shown in Plate 7, A.

Another check on the optimum temperature growth for both strains was obtained in temperature experiments on zoospore production. A suspension of spores from corn meal cultures was made in distilled and tap water. Vials containing 5 cc. of these suspensions were placed at temperatures in the Altmann incubators ranging from 7° to 32° and at room temperature exposed to the light. In no cases during these trials were zoospores produced, but an abundance of direct germination developed. The trials were run for one and three days, respectively. Very slight direct germination of the spores from either strain was noted at a temperature of 7°. Germination and growth then increased with the increase in temperature, being best between 20° and 27.5°. The optimum temperature for direct germination and growth appeared to be around 27.5°. At 30° germination and growth were not so good. The effect of light and dark at room temperatures showed no difference in the germination.

RELATION TO ACIDITY AND ALKALINITY

The relation of both organisms to acidity and alkalinity was determined by growing them on plates of potato dextrose agar and in Czapeks solution at various hydrogen-ion concentrations ranging from 1.4 to 10+ for the potato dextrose agar and from 2 to 10+ for the Czapeks solution. During the period of the experiments the room temperatures ranged between 21.5° to 27.5°, with an average temperature of approximately 23°. In both cases the vegetative growth was determined by measuring the diameter of the growth at the various concentrations. In all cases

the strains from coconut and cacao grew so nearly the same that it was impossible to note any marked difference between them.

On potato dextrose agar a distinct curve of growth for each was produced, with no growth at a hydrogen-ion concentration of 2 and below, and slight growth at 2.4. Above this concentration there was a gradual increase in the growth up to 8.8, when a decrease was noted. Between a hydrogen-ion concentration of 5.8 and 8.8 there appeared to be no marked difference in the extent of growth. Growth still took place at a concentration of 10+.

The growth between 2.4 and 3.8 was flat, dense, thick, and tough, with the production of few chlamydospores and conidia. A fluffy, dense, thick growth with few to many chlamydospores and with a smaller number of conidia was produced between 4.0 and 5.2. The mass of hyphae was still fluffy between a hydrogen-ion concentration of 5.6 and 7.8, but gradually became thinner, with a surface growth about the edges, above these concentrations. At these concentrations many chlamydospores and conidia were produced. Between 7.8 and 8.8 the aerial, fluffy growth began to disappear. Fewer spores of both kinds were formed. Above a hydrogen-ion concentration of 9 the growth was thin, flat, being on the surface of the agar, and few to no spores were formed. At 10+ it was thin on the surface, and without spores. At all concentrations except 10+ the hyphae penetrated and grew in the agar as well as upon the surface.

The optimum for growth appeared to be between a hydrogen-ion concentration of 7.4 to 7.8. Good growth was also produced up to 8.8. Slight growth took place at an acidity of 2.4. An alkalinity of 10+ did not prevent the development of either strain.

In the Czapeks solution the difference between the initial hydrogen-ion concentration and that at the end of the experiment was obtained. Control flasks of the solutions without the fungi were also kept. The growth at the different concentrations for both the coconut and cacao strains was so nearly identical that no real difference could be noted. No vegetation was obtained between hydrogen-ion concentrations of 2 and 3.2. Slight growth started at 3.4 and then gradually increased up to 9. Above this the growth gradually decreased, but still was apparent in the flasks with an initial concentration of 10+. No spores were produced in the liquid media. The optimum growth appeared to be around 7.8.

A marked change in the hydrogen-ion concentration, except in the extremely acid solutions, was noted after the fungus had grown in the media for 20 days. Between 3.4 and 3.6 the initial and final concentrations were the same. From 4.6, and above, the final concentration was less than that of the initial. The solutions with an initial concentration of from 7.8 to 10+ all were brought down to a final concentration ranging between 7 and 7.9. The controls remained the same except for a slight reduction in the concentrations above 7.8.

These tests show that both strains of the fungus were able to withstand a rather large range of concentrations from acid to alkaline. The best growth, however, appeared to be somewhat above the neutral point between 7.4 and 8.4. The minimum hydrogen-ion concentration at which growth took place was 2.4, and the maximum was 10+. Plate 7, B and C, shows the growth at the various hydrogen-ion concentrations

MORPHOLOGICAL CHARACTERISTICS

A comparative study from a morphological standpoint was made of both strains. Since no marked differences could be determined, one description only will be given to represent both. The descriptions of the morphological characters given before for the coconut strain by the writer (7) will be followed, but certain additional studies will be included.

MYCELIUM

The mycelium is white, producing a dense mass in pure culture. It contains many nuclei. In young cultures it is nonseptate and granular. In older ones septa may be produced during spore production. Protoplasmic streaming is common. The submerged mycelium (Pl. 9, A and B) is more or less gnarled, while the aerial is straight. Branching is abundant. The width varies from about 3 to about 8 μ. In the host tissue the mycelium is intercellular (Pl. 8, A). Numerous fingerlike forms of haustoria are produced, penetrating the host cells (Pl. 8, B). A distinct constriction is formed at the point of entrance into the cell. No cacao material was available at the time that the haustoria studies were made, so it cannot be stated with certainty that they are present in this strain. The similarity of both strains of the fungus in all other respects would lead one to believe that haustoria are also produced by the mycelium and present in the tissues of the cacao.

CONIDIOPHORES

Conidiophores are produced in great abundance in pure culture. Special culture methods must be employed to show them in their best condition. Material showing excellent conidiophore production may be obtained in sterilized Van Tieghem cells, by placing a few spores on a film of plain agar that has been spread over the flamed cover slip. Each conidiophore may bear from 1 to 15 or more conidia. A conidium is produced at the tip of a conidiophore; the latter then continues its growth by pushing the conidium to one side and produces another conidium at its tip. By a continuation of this process a bunch of spores is finally formed (Pl. 9, C and D). The conidiophores vary in size, ranging approximately from 180 to 645 μ in length, and from 3 to 6 μ in width.

CONIDIA

The conidia are produced terminally as described above. They are elliptical to ovate, and possess very prominent, raised, terminal papillae (Pl. 8, F and G; pl. 10, A and B). A short stalk may be present. The spores are pale yellow to colorless, are densely granular, contain many nuclei, and usually have a large vacuole (Pl. 12, A and B). The nuclei are clearly shown in prepared sections stained by means of the triple stain (Pl. 8, D, F, G). Older conidia usually possess more granules that are in groups.

MEASUREMENTS OF *conidia*.—The measurements conform closely to those of *Phytophthora faberi* Maubl., using the methods employed by Rosenbaum (*8*). Measurements were made from oatmeal agar cultures. The coconut culture was 9 to 11 days old, and that of cacao 16 to 19 days old. The lengths and widths of 400 spores were measured. The results are presented in Table I, which gives the class in microns and the number of conidia out of 400, both for length and width, that falls into each class.

TABLE I.—*Summary of measurements of conidia from coconut and cacao*

Class (in microns).	Number of coconut conidia in each class according to—		Number of cacao conidia in each class according to—	
	Length.	Width.	Length.	Width.
9.5 to 11.49	0	0	0	
11.5 to 13.49	0	0	0	
13.5 to 15.49	0	1	0	
15.5 to 17.49	0	3	0	
17.5 to 19.49	0	3	0	
19.5 to 21.49	3	6	2	
21.5 to 23.49	0	12	3	
23.5 to 25.49	2	12	2	
25.5 to 27.49	3	37	7	
27.5 to 29.49	2	32	4	
29.5 to 31.49	3	64	6	
31.5 to 33.49	13	116	12	
33.5 to 35.49	5	43	5	
35.5 to 37.49	2	27	5	
37.5 to 39.49	14	39	17	
39.5 to 41.49	13	1	4	
41.5 to 43.49	15	2	10	
43.5 to 45.49	24	2	29	
45.5 to 47.49	16	0	11	
47.5 to 49.49	24	0	17	
49.5 to 51.49	46	0	40	
51.5 to 53.49	39	0	36	
53.5 to 55.49	26	0	27	
55.5 to 57.49	36	0	40	
57.5 to 59.49	14	0	16	
59.5 to 61.49	29	0	21	
61.5 to 63.49	30	0	30	
63.5 to 65.49	10	0	12	
65.5 to 67.49	7	0	12	
67.5 to 69.49	10	0	13	
69.5 to 71.49	3	0	5	
71.5 to 73.49	3	0	3	
73.5 to 75.49	5	0	7	
75.5 to 77.49	0	0	0	
77.5 to 79.49	2	0	1	
79.5 to 81.49	3	0	1	
81.5 to 83.49	2	0	0	
83.5 to 85.49	0	0	0	
85.5 to 87.49	0	0	1	
87.5 to 89.49	0	0	1	
89.5 to 91.49	0	0	0	
Total	400	400	400	400

From Table I it can be readily seen that for the coconut strain the conidia varied in length from 19.5 to 83.49 μ and in width from 13.5 to 45.49 μ. In length the majority of the spores fell into the classes between 37.5 and 63.49 μ, the average length being 52.27 μ. In width the majority fell into classes between 19.5 and 39.49 μ, the average width being 31.28 μ. These measurements made in May, 1922, on oatmeal agar show a greater range than the measurements made in 1919 on corn-meal cultures and reported by the writer (7). The average length and width, however, remained approximately the same.

For the cacao strain the conidia varied in length from 19.5 to 89.49 μ, and in width from 11.5 to 41.49 μ. In length the majority of the spores fell into the classes between 37.5 and 63.49 μ, the average length being 51.92 μ. In width the majority fell into classes between 19.5 and 39.49 μ, the average width being 30.67 μ.

The measurements of both strains show a remarkable similarity. Figures 1 and 2 present the measurements in graphic form and clearly show the close similarity of both strains.




FIG. 1.—Graph showing the variation in length of conidia.

Table II gives the arrangement in classes of the ratios of the length to the width of the conidia for both strains.

TABLE II. *Arrangement in classes of the ratios of the length to the width of conidia, showing the limits of variation*

Class (in microns).	Number of coconut spores in each class.	Number of cacao spores in each class.
0.95 to 1.04	0	0
1.05 to 1.14	1	2
1.15 to 1.24	6	7
1.25 to 1.34	31	22
1.35 to 1.44	43	48
1.45 to 1.54	57	54
1.55 to 1.64	80	59
1.65 to 1.74	53	52
1.75 to 1.84	39	54
1.85 to 1.94	28	35
1.95 to 2.04	17	27
2.05 to 2.14	24	19
2.15 to 2.24	9	8
2.25 to 2.34	1	5
2.35 to 2.44	3	2
2.45 to 2.54	2	5
2.55 to 2.64	0	0
2.65 to 2.74	0	0
2.75 to 2.84	0	0
2.85 to 2.94	0	0
2.95 to 3.04	0	1
3.05 to 3.14	1	0
3.15 to 3.24	1	0
3.25 to 3.34	0	0
3.35 to 3.44	0	0
3.45 to 3.54	1	0
3.55 to 3.64	0	0
Total	400	400

The class of ratio values into which the greatest number of conidia fell for both strains was 1.25 to 2.14. The average ratio of length to width was 1.68 for both strains. While this ratio is somewhat larger than the measurements showed in the previous work (7), it is not of a sufficient variance, when all other morphological characters are taken into consideration, to warrant making a new species for both strains. Rosenbaum (8) in his description of *Phytophthora faberi* Maubl. gives a mean ratio of 1.47 and this figure was more clearly obtained by the writer in his previous measurements of the coconut strain (7). Graphic representations of the arrangements in classes of the ratio of the length to the width of conidia, showing the limits of variation for both strains, are given in figures 3 and 4.

GERMINATION OF THE CONIDIA.—Germination takes place by the production of either germ tubes or swarm spores. Every conidium is potentially a sporangium; its method of germination is influenced greatly by its environment. Germination by germ tubes is by far the commoner method under cultural conditions. From 1 to 5 germ tubes may be produced; these apparently may develop from any part of the surface of the conidium (Pl. 10, C). Both strains under favorable conditions, will produce an abundance of zoospores. These conditions are not clearly understood. Attempts to produce zoospores from spore suspension of corn meal cultures in sterilized and tap water by placing vials containing 5 cc. of these suspensions at various temperatures in Altmann regulated incubators from 4° up to 30° C., failed to produce zoospores. Excellent direct germination was obtained. Frequently old oatmeal agar cultures, from which the surface mycelium had been removed and then new growth allowed to start, would produce zoospores by the addition of water. Swarm spores were also obtained in Van Tieghem cells by growing the organisms on very dilute agar or in hanging drops of water placed on sterile cover slips. Just before formation there appeared a rearrangement of the protoplasmic granules. The swarm spores were then produced within the sporangium (Pl. 11, A and B). The end of the papilla finally broke off or was dissolved and the spores escaped. No vesicle formation has been observed. The spores approach the opening and escape one by one. When produced on dilute agar the zoospores oozed out of the sporangium and remained in a mass at the mouth. Here they soon germinated by sending out germ tubes (Pl. 11, C and D). Frequently zoospores did not escape from the sporangium and then germination took place within the sporangium (Pl. 11, C and D).




FIG. 2.—Graph showing the variation in width of conidia.

The process of emerging from the sporangium in water cultures is extremely interesting. After the dissolution of the tip of the papilla an active movement of the developed zoospores within takes place. As

soon as a zoospore finds the opening it comes to rest and gradually oozes out, becoming much elongated in the process and constricted in the middle and bulging out at both ends, due to the smallness of the opening (Pl. 11, A and B). After the zoospore passes through, it comes to rest for a portion of a second and then swims off actively. While one zoospore is passing out, the rest within swim about against the wall until the opening is again free, when one by one they will escape in the same manner. The zoospores are granular, slightly greenish, and have distinct nuclei and one or two vacuoles. From a side view they are kidney shaped, and when observed from the end a groove is seen on one side from tip to tip. The swarm spores swim about for a time by means of two flagella. Just




FIG. 3.—Arrangement in classes of the ratios of the length to the width of conidia (coconut), showing the limits of variation.

before coming to a permanent rest the spores become sluggish, settle down for a time, then start off, settle down again, and swim away once more, and finally settle down again definitely. No distinct spore wall was noted on the motile spores, but as soon as they came to a permanent rest a wall could be distinguished, which was especially clear when germination started (Pl. 11, A and B). In the rest stage the spores became spherical, in some cases with a depression on one side, took on a more greenish tinge, became more granular, enlarged somewhat, and then put out germ tubes which grew into a mycelium (Pl. 11, A and B). In 25 minutes the germ tube of one zoospore on potato dextrose agar was observed to double in length. Frequently in hanging drop slides of water and dilute agar all the zoospores were produced from secondary sporangia that arose from the original spores placed in the drop or on the agar.

The original spore in these cases first germinated by the production of a germ tube that grew into a short mycelium, which in turn became a sporangiophore producing a group of sporangia. Some of the latter would then produce zoospores.

CHLAMYDOSPORES

Chlamydospores are produced directly from the mycelium, usually terminally, but sometimes intercalarily (Pl. 12, C and D). They are spherical, granular like the conidia, but with a slightly deeper yellow shade (Pl. 8, E). In pure culture they are produced in great abundance on oatmeal or potato dextrose agar. In the diseased rind of cacao fruits many thick-walled chlamydospores are usually produced (Pl. 8, F).




FIG. 4.—Arrangement in classes of the ratios of the length to the width of conidia (cacao), showing the limits of variation.

MEASUREMENTS OF CHLAMYDOSPORES.—The size of the chlamydospores is one of the criteria by which the species of Phytophthora are separated. Four hundred measurements were made of the diameters of the spores from both coconut and cacao strains grown on oatmeal agar. The coconut culture was 9 to 10 days old, and that of cacao was 11 days old. A summary of these measurements, grouped in classes, appears in Table III.

The chlamydospores of the coconut strain varied in diameter from 19.5 to 61.49 μ, the average diameter being 41.62 μ. These measurements were made in May, 1922, of spores grown on oatmeal agar and correspond favorably with those made in 1919 grown on corn meal (7).

For the cacao strain the chlamydospores varied in diameter from 17.5 to 53.49 μ, the average diameter being 41.06 μ.

TABLE III.—*Summary of measurements of chlamydospores from coconut and cacao*

Class (in microns).	Number of chlamydospores in each class from—	
	Coconut.	Cacao.
15.5 to 17.49	0	0
17.5 to 19.49	0	1
19.5 to 21.49	4	0
21.5 to 23.49	1	0
23.5 to 25.49	1	1
25.5 to 27.49	3	3
27.5 to 29.49	5	3
29.5 to 31.49	1	4
31.5 to 33.49	10	7
33.5 to 35.49	6	8
35.5 to 37.49	16	25
37.5 to 39.49	70	69
39.5 to 41.49	49	69
41.5 to 43.49	69	81
43.5 to 45.49	91	99
45.5 to 47.49	37	17
47.5 to 49.49	22	8
49.5 to 51.49	12	3
51.5 to 53.49	1	2
53.5 to 55.49	1	0
55.5 to 57.49	0	0
57.5 to 59.49	0	0
59.5 to 61.49	1	0
61.5 to 63.49	0	0
Total	400	400

The measurements, therefore, for both strains are so nearly identical that from this standpoint both fall under the same species. A graphic representation of the measurements for both strains is given in figure 5.

GERMINATION OF CHLAMYDOSPORES.—In Van Tieghem cells, prepared with hanging drops of distilled water, corn meal extract, potato dextrose agar, or pure agar, direct germination took place within 24 hours. From 1 to 12 germ tubes may arise from one spore.

SEXUAL BODIES

No sexual bodies have been observed in diseased portions of coconut or cacao trees or in pure cultures of either of these strains. The absence of antheridia places these strains in the Faberi group, in accordance with Rosenbaum (*8*), "which embraces the forms in which the antheridia are entirely absent or in which the relation of the antheridium to the oogonium is unknown."

TAXONOMY

According to the tentative table offered for the separation of species of Phytophthora and devised by Rosenbaum (*8*), both strains fall into the Faberi group and in the species *Phytophthora faberi* Maubl. The discrepancy in the case of the ratio of the length to the width of the conidia, alone, does not warrant the creation of a new species for either strain, especially in the present state of the Phytophthora group. The descriptions and measurements of both strains also correspond to those of the original description of the species by Maublanc (*4, p. 314–324*).

DISCUSSION

RELATIONSHIP

The two strains of Phytophthora are, therefore, to be considered as identical. Both behave alike in infection experiments. The physiological characteristics of each from a standpoint of growth on various media, relation to light, relation to temperature, and relation to acidity and alkalinity are the same. The morphological characteristics are also so nearly alike that no real distinction can be made between them. Both strains of the fungus correspond closely to the Faberi group and to the species *Phytophthora faberi* Maubl. as determined by Rosenbaum (*8*). Antheridia are entirely unknown. The papillae of the conidia are raised and very prominent. The measurements of conidia and chlamydospores closely approximate those for this species. There is a slight variation in the ratio of the length to the width of conidia. This difference, when taken into consideration with all the other points of similarity, is not sufficient to warrant the creation of a new species. Rosenbaum (*8*) gives 1.47 as the mean ratio of length to width of conidia of *Phytophthora faberi* Maubl. The ratio obtained in the last comparative measurements for both strains on oatmeal agar was 1.60. Measurements made in 1919 (*7*) of the coconut strain grown on corn meal gave a mean ratio of 1.51, being more nearly that as given by Rosenbaum. The slight change in the form of the spores measured in 1919 and 1922 can not be accounted for unless it be due to a difference in age and a change in the medium used.




FIG. 5.—Graph showing the variation in the diameter of chlamydospores.

The original description of *Phytophthora faberi* Maubl. as given by Maublanc (*4*) in 1909 on cacao (*Theobroma cacao*) corresponds also very closely to that of the two strains studied by the writer.

In 1907 Butler (*2*) described a new species of a Phycomycete occurring on various hosts, among which was the coconut (*Cocos nucifera*), under the name of *Pythium palmivorum*. Later in 1918 (*3*), the genus was transferred to that of Phytophthora without redescription. In 1907 Ashby (*1*), in an article on two diseases of coconut palm in Jamaica, ascribed the cause of the budrot to *Phytophthora palmivorum* Butl. after having submitted a culture to Butler for comparison. A culture of the fungus from Jamaica (Ashby, subculture of 31-8-20), obtained from Mr. A. Sharples, mycologist of the Federated Malay States, appears to be very much like that of the Phytophthora on coconut and cacao from the Philippines. No accurate comparisons, however, were made. The Phytophthora from the Philippines in all probability is different from that in India, as originally described under *Pythium*

palmivorum, and then later referred to *Phytophthora palmivorum*, as the latter was reported to produce antheridia and oogonia. The development of these organs was not, however, determined with certainty. Ashby (*1*) also observed antheridia and oogonia in his strain from coconut.

In addition to these two hosts, Phytophthoras have been obtained from the following various plants in the Philippines: Two apparently different strains causing the fruitrot and blight of eggplant (*Solanum melongena*); a strain causing the blight of roselle seedlings (*Hibiscus sabdariffa*); a strain causing the blight of citrus seedlings and buds of budded plants (*Citrus* spp.); a strain producing the rot of abaca suckers (*Musa textilis*); and a strain causing the blight of Hevea rubber (*Hevea brasiliensis*). From a general observation some of these strains appear to be identical with those described from coconut and cacao. However, until a complete comparative test is made, no definite statement as to their identity can be given. The determination of the relationship of these various strains is of great importance in a study of tropical diseases of plants.

SIGNIFICANCE OF SIMILARITY OF BOTH STRAINS

The discovery of the similarity of the strains from coconut and cacao is of extreme consequence from a control standpoint. In certain coconut sections it is a common practice to interplant with cacao. As the latter tree is practically always infested with the Phytophthora, this custom will have to be discontinued. It seems highly probable that a number of the other strains listed above may be identical with those on coconut and cacao. A study of these would prove to be of great interest and would bring out new means of combating the diseases.

SUMMARY

Coconut culture is one of the chief industries of the Philippine Islands. Millions of trees have been planted on the various islands of the group. The infectious type of budrot has made its appearance in certain sections, primarily in the extensive coconut regions in Laguna, Batangas, and Tayabas Provinces on the island of Luzon. Here sporadic epidemics of the disease have been authentically reported since 1908, and thousands of dollars of loss has occurred since that time.

Various organisms, including bacteria and fungi, have been considered responsible for the disease. In the Philippine Islands there are apparently two types of budrot, the really infectious type caused by a Phytophthora, and a secondary type following some injury such as beetle injury and caused primarily by the invasion of bacteria in the weakened tissue.

The culture of cacao in the Philippine Islands, while not extensive, can be considered as one of the secondary industries. The manner in which the trees are grown by the average farmer subjects the trees to severe fungus attacks. In certain sections frequently one-half of the cacao crop is destroyed. The canker of the branches and trunk, and the rot of the pods are the chief troubles, and they are caused by a Phytophthora. Cacao trees are often interplanted with coconut trees, offering a ready means of transfer of a fungus from one plant to the other.

A careful study of the cross infection possibilities and the physiological and morphological characteristics of the fungus strains from the

coconut and cacao has shown that the coconut budrot and the blackrot and canker of cacao are caused by the same species of Phycomycete, *Phytophthora faberi* Maubl. The discovery of the identity of these two strains is of utmost importance from a control standpoint. Coconut and cacao trees should not be interplanted, since the fungus can be readily transmitted from one host to the other.

Similar types of coconut budrot in India and Jamaica have been shown by the investigators in those countries to be due to a Phytophthora. The similarity of the strains on coconut and cacao has been determined only in the Philippine Islands. A comparative study of the Phytophthoras from India and Jamaica would be of interest.

The presence of other strains of Phytophthora on important economic plants in the Philippine Islands, with the possibilities of cross inoculation between these, indicates that some may be identical with the strains on coconut and cacao.

LITERATURE CITED

(1) ASHBY, S. F.
1920. NOTES ON TWO DISEASES OF THE COCO-NUT PALM IN JAMAICA CAUSED BY FUNGI OF THE GENUS PHYTOPHTHORA. *In* West Indian Bul., v. 18, p. 61–73.

(2) BUTLER, E. J.
1907. AN ACCOUNT OF THE GENUS PYTHIUM AND SOME CHYTRIDIACEAE. *In* Mem. Dept. Agr. India Bot. Ser., v. 1, no. 5, 160 p., 10 pl. Literature, p. 142–147.

(3) ———
1918. FUNGI AND DISEASE IN PLANTS. iv, 547 p., 206 fig. Calcutta. Bibliography, p. 518–531.

(4) MAUBLANC, André.
1909–10. LES MALADIES DES PLANTES CULTIVÉES DANS LES PAYS CHAUDS. Maladies du cacaoyer. *In* Agr. Prat. Pays Chauds, ann. 9, p. 312–324, 393–407, 472–479; 10, p. 510–512, illus.

(5) PETHYBRIDGE, George H.
1913. ON THE ROTTING OF POTATO TUBERS BY A NEW SPECIES OF PHYTOPHTHORA HAVING A METHOD OF SEXUAL REPRODUCTION HITHERTO UNDESCRIBED. *In* Sci. Proc. Roy. Dublin Soc., n. s., v. 13, p. 529–564, pl. 42–44. Bibliography, p. 561–562.

(6) REINKING, Otto A.
1918. PHILIPPINE ECONOMIC-PLANT DISEASES. *In* Philippine Jour. Sci. Sec. A., v. 13, p. 165–216, 217–274, 43 fig., 22 pl.


(7) ———
1919. PHYTOPHTHORA FABERI MAUBL.: THE CAUSE OF COCONUT BUD ROT IN THE PHILIPPINES. *In* Philippine Jour. Sci., v. 14, p. 131–151, 3 pl. Bibliography, p. 149–150.


(8) ROSENBAUM, J.
1917. STUDIES OF THE GENUS PHYTOPHTHORA. *In* Jour. Agr. Research, v. 8, p. 233–276, 13 fig., pl. 71–77. Literature cited, p. 273–276.

PLATE 1

A.—Typical field case of coconut budrot showing infection of region about growing point and in bud. From this specimen, obtained on March 11, 1919, at Lilio, Laguna Province, P. I., the pure culture of the fungus strain used for the coconut experiments was obtained. The photomicrographs (Pl. 8) showing the mycelium and haustoria in the tissue were made from sections obtained from this particular specimen.

B.—Early stages in rot of papaya (*Carica papaya*) produced by inoculation with Phytophthora strain from coconut. A similar infection is produced by the cacao strain.


B
A
USDA


B



A

PLATE 2

Canker of cacao branches and blackrot of cacao pods.

A.—Canker of cacao branches produced by inoculation with Phytophthora strain from cacao.

B. Blackrot of cacao pod produced by inoculation with Phytophthora strain from cacao.

PLATE 3

Typical cases of coconut budrot produced by inoculation with Phytophthora strains from coconut and cacao.

A.—Budrot produced in healthy, vigorously growing coconut tree by merely pouring a zoospore suspension in rain water of the coconut strain between the growing leaves of the bud. Bud killed in 15 to 16 months after inoculation.

B.—Budrot produced in a disease-free coconut tree by inoculation with a zoospore suspension of the cacao strain. The tree was also infested with beetles, as indicated by the leaves with triangular shaped portions cut out of the leafleats. Bud killed 2 to 3 months after inoculation. The rapid production of the rot in this case was undoubtedly due to the beetle injury to the leaves, and also to the fact that the tree was not in a vigorous condition of growth.


B




A
USDA
B

PLATE 4


Coconut budrot produced by inoculation of healthy tree with Phytophthora strain from coconut.

A.—Longitudinal section of diseased tree shown in Plate 3, A, showing typical rot of region about growing point and in lower part of bud.


B.—Diseased tree, on right, and healthy control tree, on left, 15 to 16 months after inoculation. At time of inoculation both trees were of approximately the same age and height. The stunting of the infected tree is clearly shown.

PLATE 5

Coconut seedlings inoculated with Phytophthora strain from coconut.
A.—Inoculated tree and control. Stab method of inoculation employed.
B.—Typical and severe rot produced in seedling, using stab method of inoculation.



B



A



A
B
C
D
E

PLATE 6

Rot of tomato, potato, and apple produced by inoculation with both strains of Phytophthora.

A.—Rot of ripe tomato produced by inoculation with Phytophthora strain from cacao.

B.—Rot of ripe tomato produced by inoculation with Phytophthora strain from coconut.

C.—Rot of potato produced by inoculation with Phytophthora strain from coconut. Rotted area pinkish in color. A similar rot is produced by Phytophthora strain from cacao.

D.—Rot of Ben Davis apple produced by inoculation with Phytophthora strain from cacao.


E.—Rot of Ben Davis apple produced by inoculation with Phytophthora strain from coconut.

PLATE 7

A.—Growth of coconut strains on potato dextrose agar made in Altman incubators at various temperatures. The effect of light and darkness is clearly shown in the two plates grown at the same temperature: 22.0 in the dark; 22.0L, in the light. A fluffy granular growth due to sporulation is produced in the light.

B.—Growth on potato dextrose agar at room temperature of the Phytophthora strain from coconut under various hydrogen-ion concentrations.

C.—Growth on potato dextrose agar at room temperature of the Phytophthora strain from cacao under various hydrogen-ion concentrations.


A
B
C



A
B
C
D
E
F
G

PLATE 8

Photomicrographs of diseased tissues of coconut and cacao:

A.—Section from diseased tissue near growing point of field case of coconut budrot, showing intercellular, nonseptate mycelium. Approximately, × 200. Acid fuchsin and light-green stain.

B.—Section from diseased tissue near growing point of field case of coconut budrot, showing fingerlike haustoria penetrating host cells. Approximately, × 750. Acid fuchsin and light-green stain.

C.—Free-hand section of rind of diseased cacao pod, showing chlamydospores. Approximately, × 250. Acid fuchsin and light-green stain.

Photomicrographs of conidia and chlamydospores of both strains of Phytophthora:

D.—Conidia of Phytophthora strain from coconut, showing presence of many nuclei. Sectioned material from pure culture of fungus on oatmeal agar. Approximately, × 750. Triple stain.

E.—Chlamydospores of Phytophthora strain from cacao. Sectioned material from pure culture on oatmeal agar. Approximately, × 750. Triple stain.

F.—Conidium of Phytophthora strain from coconut, showing papilla stained deeper than wall of spore, and numerous nuclei. Sectioned material from pure culture on oatmeal agar. Approximately, × 750. Triple stain.


G.—Conidia of Phytophthora strain from cacao showing presence of many nuclei. Sectioned material from pure culture of fungus on oatmeal agar. Approximately, × 750. Triple stain.


PLATE 9


Mycelium, conidiophores, and conidia of both strains of Phytophthora.

A, B.—Submerged mycelium of both strains of Phytophthora grown on potato agar. × 400. A, coconut. B, cacao.

C, D.—Conidiophores and conidia of both strains of Phytophthora. × 400. C, coconut. D, cacao.


C
D


A
B


C



A
B

PLATE 10

A, B.—Prominent, raised, terminal papillae from conidia of both strains of Phytophthora grown on potato agar. × 400. A, coconut. B, cacao.

C.—Direct germination of conidia of the Phytophthora strain from coconut showing the production of from few to many germ tubes. × 400.


54484—23——4


PLATE 11


Zoospore production and germination in various stages of development of both stages of Phytophthora.

A, B.—Zoospore production showing various stages of development and germination. × 400. A, coconut. B, cacao.

C, D.—Zoospore germination in place within the sporangium and at the mouth of the sporangium. × 400. Produced in Van Tieghem cells on dilute agar. C, coconut. B, cacao.


C
D
USDA


A
B
USDA


C
D



B
A

PLATE 12

Various types of conidia and chlamydospores of both strains of Phytophthora.
A, B.—Conidia of various forms of both strains of Phytophthora. × 400. A, coconut. B, cacao.
C, D.—Chlamydospores of various sizes of both strains of Phytophthora. × 400. C, coconut. D, cacao.